

Table 3.1C Perceived Risk and Availability of Drugs, by Age Group: Standard Errors of Numbers in Thousands, 2002

Risk/Availability ¹	Total	AGE GROUP (Years)		
		12-17	18-25	26 or Older
PERCEPTIONS OF GREAT RISK				
Cigarettes				
Smoke one or more packs per day	750	108	138	732
Marijuana				
Smoke once a month	912	97	130	872
Smoke once or twice a week	881	105	148	828
Cocaine				
Use once a month	757	108	140	726
Use once or twice a week	484	86	86	477
Heroin				
Try once or twice	563	108	111	536
Use once or twice a week	363	82	73	342
LSD				
Try once or twice	696	109	140	670
Use once or twice a week	474	93	92	464
Alcohol				
Four or five drinks nearly every day	775	118	139	764
Five or more drinks once or twice a week	863	106	145	834
PERCEIVED AVAILABILITY				
Fairly or very easy to obtain				
Marijuana	898	109	131	873
Cocaine	816	86	136	792
Crack	821	90	133	793
Heroin	669	73	110	649
LSD	661	79	117	638
Approached by someone selling drugs in the past month	359	82	110	330

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 3.1D Perceived Risk and Availability of Drugs, by Age Group: Standard Errors of Percentages, 2002

Risk/Availability ¹	Total	AGE GROUP (Years)		
		12-17	18-25	26 or Older
PERCEPTIONS OF GREAT RISK				
Cigarettes				
Smoke one or more packs per day	0.32	0.44	0.45	0.41
Marijuana				
Smoke once a month	0.39	0.40	0.42	0.50
Smoke once or twice a week	0.38	0.43	0.48	0.47
Cocaine				
Use once a month	0.33	0.44	0.45	0.41
Use once or twice a week	0.21	0.35	0.28	0.27
Heroin				
Try once or twice	0.24	0.45	0.36	0.30
Use once or twice a week	0.16	0.34	0.24	0.19
LSD				
Try once or twice	0.30	0.46	0.46	0.38
Use once or twice a week	0.21	0.39	0.30	0.26
Alcohol				
Four or five drinks nearly every day	0.33	0.48	0.45	0.43
Five or more drinks once or twice a week	0.37	0.43	0.47	0.47
PERCEIVED AVAILABILITY				
Fairly or very easy to obtain				
Marijuana	0.40	0.45	0.43	0.51
Cocaine	0.37	0.36	0.45	0.47
Crack	0.37	0.37	0.44	0.47
Heroin	0.30	0.31	0.37	0.39
LSD	0.30	0.34	0.39	0.38
Approached by someone selling drugs in the past month	0.15	0.33	0.35	0.18

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 3.2C Perceived Great Risk of Smoking One or More Packs of Cigarettes Per Day, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Smoking One or More Packs of Cigarettes Per Day (II) among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002

(I) Demographic Characteristic	Smoke One or More Packs of Cigarettes Per Day ¹	(II) Drug Use	Smoke One or More Packs of Cigarettes Per Day ¹	
	Great Risk		Great Risk	Moderate/Slight/No Risk
TOTAL	108	PAST MONTH		
AGE		Any Illicit Drug ²	52	46
12-13	62	Marijuana	43	39
14-15	62	Any Illicit Drug Other Than Marijuana ²	38	34
16-17	53	Cigarettes	51	50
GENDER		Alcohol	64	51
Male	76	Binge Alcohol ³	47	46
Female	69	PAST YEAR		
HISPANIC ORIGIN AND RACE		Any Illicit Drug ²	69	59
Not Hispanic or Latino	97	Marijuana	58	53
White	80	Any Illicit Drug Other Than Marijuana ²	56	48
Black or African American	35	Cigarettes	64	57
American Indian or Alaska Native	7	Alcohol	83	62
Native Hawaiian or Other Pacific Islander	*	LIFETIME		
Asian	26	Any Illicit Drug ²	76	66
Two or More Races	11	Marijuana	65	59
Hispanic or Latino	49	Any Illicit Drug Other Than Marijuana ²	66	57
COUNTY TYPE		Cigarettes	84	61
Large Metro	83	Alcohol	82	65
Small Metro	55			
Nonmetro	42			

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 3.2D Perceived Great Risk of Smoking One or More Packs of Cigarettes Per Day, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Smoking One or More Packs of Cigarettes Per Day (II) among Persons Aged 12 to 17: Standard Errors of Percentages, 2002

(I) Demographic Characteristic	Smoke One or More Packs of Cigarettes Per Day ¹	(II) Drug Use	Smoke One or More Packs of Cigarettes Per Day ¹	
	Great Risk		Great Risk	Moderate/Slight/No Risk
TOTAL	0.44	PAST MONTH		
AGE		Any Illicit Drug ²	0.34	0.51
12-13	0.76	Marijuana	0.28	0.43
14-15	0.74	Any Illicit Drug Other Than Marijuana ²	0.25	0.38
16-17	0.67	Cigarettes	0.33	0.55
GENDER		Alcohol	0.41	0.56
Male	0.61	Binge Alcohol ³	0.30	0.51
Female	0.57	PAST YEAR		
HISPANIC ORIGIN AND RACE		Any Illicit Drug ²	0.45	0.65
Not Hispanic or Latino	0.47	Marijuana	0.37	0.58
White	0.52	Any Illicit Drug Other Than Marijuana ²	0.36	0.53
Black or African American	0.99	Cigarettes	0.42	0.63
American Indian or Alaska Native	3.94	Alcohol	0.53	0.68
Native Hawaiian or Other Pacific Islander	*	LIFETIME		
Asian	2.64	Any Illicit Drug ²	0.49	0.73
Two or More Races	2.94	Marijuana	0.42	0.65
Hispanic or Latino	1.28	Any Illicit Drug Other Than Marijuana ²	0.42	0.63
COUNTY TYPE		Cigarettes	0.54	0.68
Large Metro	0.67	Alcohol	0.53	0.71
Small Metro	0.75			
Nonmetro	0.84			

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 3.3C Perceived Great Risk of Smoking Marijuana Once a Month, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Smoking Marijuana Once a Month (II) among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002

(I) Demographic Characteristic	Smoke Marijuana Once a Month ¹	(II) Drug Use	Smoke Marijuana Once a Month ¹	
	Great Risk		Great Risk	Moderate/Slight/No Risk
TOTAL	97	PAST MONTH		
AGE		Any Illicit Drug ²	24	66
12-13	58	Marijuana	16	55
14-15	58	Any Illicit Drug Other Than Marijuana ²	21	46
16-17	53	Cigarettes	29	66
GENDER		Alcohol	34	72
Male	64	Binge Alcohol ³	24	61
Female	70	PAST YEAR		
HISPANIC ORIGIN AND RACE		Any Illicit Drug ²	34	85
Not Hispanic or Latino	89	Marijuana	24	73
White	77	Any Illicit Drug Other Than Marijuana ²	29	68
Black or African American	37	Cigarettes	38	77
American Indian or Alaska Native	7	Alcohol	48	89
Native Hawaiian or Other Pacific Islander	*	LIFETIME		
Asian	25	Any Illicit Drug ²	44	92
Two or More Races	11	Marijuana	30	79
Hispanic or Latino	41	Any Illicit Drug Other Than Marijuana ²	39	79
COUNTY TYPE		Cigarettes	51	88
Large Metro	71	Alcohol	56	89
Small Metro	49			
Nonmetro	39			

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 3.3D Perceived Great Risk of Smoking Marijuana Once a Month, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Smoking Marijuana Once a Month (II) among Persons Aged 12 to 17: Standard Errors of Percentages, 2002

(I) Demographic Characteristic	Smoke Marijuana Once a Month ¹	(II) Drug Use	Smoke Marijuana Once a Month ¹	
	Great Risk		Great Risk	Moderate/Slight/No Risk
TOTAL	0.40	PAST MONTH		
AGE		Any Illicit Drug ²	0.30	0.40
12-13	0.72	Marijuana	0.20	0.34
14-15	0.69	Any Illicit Drug Other Than Marijuana ²	0.26	0.28
16-17	0.67	Cigarettes	0.37	0.40
GENDER		Alcohol	0.43	0.44
Male	0.51	Binge Alcohol ³	0.31	0.37
Female	0.59	PAST YEAR		
HISPANIC ORIGIN AND RACE		Any Illicit Drug ²	0.42	0.52
Not Hispanic or Latino	0.43	Marijuana	0.30	0.44
White	0.50	Any Illicit Drug Other Than Marijuana ²	0.37	0.41
Black or African American	1.03	Cigarettes	0.47	0.46
American Indian or Alaska Native	3.92	Alcohol	0.60	0.54
Native Hawaiian or Other Pacific Islander	*	LIFETIME		
Asian	2.54	Any Illicit Drug ²	0.56	0.56
Two or More Races	2.79	Marijuana	0.38	0.48
Hispanic or Latino	1.07	Any Illicit Drug Other Than Marijuana ²	0.49	0.48
COUNTY TYPE		Cigarettes	0.64	0.53
Large Metro	0.58	Alcohol	0.71	0.54
Small Metro	0.68			
Nonmetro	0.80			

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 3.4C Perceived Great Risk of Smoking Marijuana Once or Twice a Week, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Smoking Marijuana Once or Twice a Week (II) among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002

(I) Demographic Characteristic	Smoke Marijuana Once or Twice a Week ¹	(II) Drug Use	Smoke Marijuana Once or Twice a Week ¹	
	Great Risk		Great Risk	Moderate/Slight/No Risk
TOTAL	105	PAST MONTH		
AGE		Any Illicit Drug ²	30	59
12-13	63	Marijuana	19	53
14-15	60	Any Illicit Drug Other Than Marijuana ²	26	42
16-17	59	Cigarettes	36	60
GENDER		Alcohol	42	64
Male	72	Binge Alcohol ³	30	58
Female	72	PAST YEAR		
HISPANIC ORIGIN AND RACE		Any Illicit Drug ²	46	75
Not Hispanic or Latino	94	Marijuana	34	67
White	81	Any Illicit Drug Other Than Marijuana ²	38	61
Black or African American	38	Cigarettes	47	69
American Indian or Alaska Native	8	Alcohol	63	72
Native Hawaiian or Other Pacific Islander	*	LIFETIME		
Asian	26	Any Illicit Drug ²	57	77
Two or More Races	11	Marijuana	42	71
Hispanic or Latino	48	Any Illicit Drug Other Than Marijuana ²	50	69
COUNTY TYPE		Cigarettes	67	76
Large Metro	78	Alcohol	72	70
Small Metro	54			
Nonmetro	40			

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 3.4D Perceived Great Risk of Smoking Marijuana Once or Twice a Week, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Smoking Marijuana Once or Twice a Week (II) among Persons Aged 12 to 17: Standard Errors of Percentages, 2002

(I) Demographic Characteristic	Smoke Marijuana Once or Twice a Week ¹	(II) Drug Use	Smoke Marijuana Once or Twice a Week ¹	
	Great Risk		Great Risk	Moderate/Slight/No Risk
TOTAL	0.43	PAST MONTH		
AGE		Any Illicit Drug ²	0.24	0.50
12-13	0.78	Marijuana	0.15	0.45
14-15	0.71	Any Illicit Drug Other Than Marijuana ²	0.21	0.35
16-17	0.74	Cigarettes	0.29	0.51
GENDER		Alcohol	0.33	0.54
Male	0.58	Binge Alcohol ³	0.24	0.49
Female	0.60	PAST YEAR		
HISPANIC ORIGIN AND RACE		Any Illicit Drug ²	0.37	0.63
Not Hispanic or Latino	0.46	Marijuana	0.27	0.57
White	0.53	Any Illicit Drug Other Than Marijuana ²	0.30	0.52
Black or African American	1.06	Cigarettes	0.37	0.58
American Indian or Alaska Native	4.52	Alcohol	0.50	0.61
Native Hawaiian or Other Pacific Islander	*	LIFETIME		
Asian	2.68	Any Illicit Drug ²	0.45	0.65
Two or More Races	2.97	Marijuana	0.33	0.59
Hispanic or Latino	1.24	Any Illicit Drug Other Than Marijuana ²	0.40	0.58
COUNTY TYPE		Cigarettes	0.53	0.64
Large Metro	0.64	Alcohol	0.57	0.59
Small Metro	0.74			
Nonmetro	0.82			

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 3.5C Perceived Great Risk of Using Cocaine Once a Month, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Using Cocaine Once a Month (II) among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002

(I) Demographic Characteristic	Use Cocaine Once a Month ¹	(II) Drug Use	Use Cocaine Once a Month ¹	
	Great Risk		Great Risk	Moderate/Slight/No Risk
TOTAL	108	PAST MONTH		
AGE		Any Illicit Drug ²	44	53
12-13	60	Marijuana	39	43
14-15	62	Any Illicit Drug Other Than Marijuana ²	30	42
16-17	59	Cigarettes	52	52
GENDER		Alcohol	57	57
Male	75	Binge Alcohol ³	46	47
Female	73	PAST YEAR		
HISPANIC ORIGIN AND RACE		Any Illicit Drug ²	59	69
Not Hispanic or Latino	99	Marijuana	55	56
White	80	Any Illicit Drug Other Than Marijuana ²	46	58
Black or African American	39	Cigarettes	63	62
American Indian or Alaska Native	10	Alcohol	74	74
Native Hawaiian or Other Pacific Islander	*	LIFETIME		
Asian	25	Any Illicit Drug ²	66	77
Two or More Races	12	Marijuana	61	61
Hispanic or Latino	49	Any Illicit Drug Other Than Marijuana ²	58	68
COUNTY TYPE		Cigarettes	74	73
Large Metro	81	Alcohol	78	80
Small Metro	55			
Nonmetro	46			

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 3.5D Perceived Great Risk of Using Cocaine Once a Month, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Using Cocaine Once a Month (II) among Persons Aged 12 to 17: Standard Errors of Percentages, 2002

(I) Demographic Characteristic	Use Cocaine Once a Month ¹	(II) Drug Use	Use Cocaine Once a Month ¹	
	Great Risk		Great Risk	Moderate/Slight/No Risk
TOTAL	0.44	PAST MONTH		
AGE		Any Illicit Drug ²	0.36	0.44
12-13	0.75	Marijuana	0.32	0.35
14-15	0.75	Any Illicit Drug Other Than Marijuana ²	0.25	0.34
16-17	0.74	Cigarettes	0.43	0.43
GENDER		Alcohol	0.46	0.47
Male	0.61	Binge Alcohol ³	0.37	0.39
Female	0.61	PAST YEAR		
HISPANIC ORIGIN AND RACE		Any Illicit Drug ²	0.48	0.57
Not Hispanic or Latino	0.48	Marijuana	0.45	0.46
White	0.52	Any Illicit Drug Other Than Marijuana ²	0.38	0.48
Black or African American	1.09	Cigarettes	0.52	0.51
American Indian or Alaska Native	5.26	Alcohol	0.60	0.62
Native Hawaiian or Other Pacific Islander	*	LIFETIME		
Asian	2.58	Any Illicit Drug ²	0.54	0.64
Two or More Races	3.16	Marijuana	0.49	0.51
Hispanic or Latino	1.26	Any Illicit Drug Other Than Marijuana ²	0.47	0.56
COUNTY TYPE		Cigarettes	0.60	0.61
Large Metro	0.66	Alcohol	0.63	0.66
Small Metro	0.76			
Nonmetro	0.95			

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 3.6C Perceived Great Risk of Using Cocaine Once or Twice a Week, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Using Cocaine Once or Twice a Week (II) among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002

(I) Demographic Characteristic	Use Cocaine Once or Twice a Week ¹	(II) Drug Use	Use Cocaine Once or Twice a Week ¹	
	Great Risk		Great Risk	Moderate/Slight/No Risk
TOTAL	86	PAST MONTH		
AGE		Any Illicit Drug ²	58	37
12-13	60	Marijuana	50	29
14-15	49	Any Illicit Drug Other Than Marijuana ²	40	30
16-17	43	Cigarettes	64	34
GENDER		Alcohol	69	37
Male	60	Binge Alcohol ³	58	32
Female	60	PAST YEAR		
HISPANIC ORIGIN AND RACE		Any Illicit Drug ²	76	46
Not Hispanic or Latino	76	Marijuana	68	36
White	58	Any Illicit Drug Other Than Marijuana ²	60	40
Black or African American	34	Cigarettes	78	40
American Indian or Alaska Native	9	Alcohol	89	49
Native Hawaiian or Other Pacific Islander	*	LIFETIME		
Asian	26	Any Illicit Drug ²	84	49
Two or More Races	11	Marijuana	76	39
Hispanic or Latino	39	Any Illicit Drug Other Than Marijuana ²	71	47
COUNTY TYPE		Cigarettes	95	46
Large Metro	63	Alcohol	93	49
Small Metro	46			
Nonmetro	33			

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 3.6D Perceived Great Risk of Using Cocaine Once or Twice a Week, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Using Cocaine Once or Twice a Week (II) among Persons Aged 12 to 17: Standard Errors of Percentages, 2002

(I) Demographic Characteristic	Use Cocaine Once or Twice a Week ¹	(II) Drug Use	Use Cocaine Once or Twice a Week ¹	
	Great Risk		Great Risk	Moderate/Slight/No Risk
TOTAL	0.35	PAST MONTH		
AGE		Any Illicit Drug ²	0.30	0.74
12-13	0.75	Marijuana	0.26	0.58
14-15	0.58	Any Illicit Drug Other Than Marijuana ²	0.21	0.61
16-17	0.55	Cigarettes	0.33	0.69
GENDER		Alcohol	0.35	0.75
Male	0.48	Binge Alcohol ³	0.30	0.65
Female	0.50	PAST YEAR		
HISPANIC ORIGIN AND RACE		Any Illicit Drug ²	0.39	0.93
Not Hispanic or Latino	0.37	Marijuana	0.35	0.73
White	0.38	Any Illicit Drug Other Than Marijuana ²	0.31	0.82
Black or African American	0.95	Cigarettes	0.40	0.82
American Indian or Alaska Native	4.83	Alcohol	0.46	1.00
Native Hawaiian or Other Pacific Islander	*	LIFETIME		
Asian	2.61	Any Illicit Drug ²	0.43	1.00
Two or More Races	2.78	Marijuana	0.39	0.79
Hispanic or Latino	1.01	Any Illicit Drug Other Than Marijuana ²	0.37	0.95
COUNTY TYPE		Cigarettes	0.49	0.93
Large Metro	0.52	Alcohol	0.48	0.99
Small Metro	0.64			
Nonmetro	0.68			

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 3.7C Perceived Great Risk of Trying Heroin Once or Twice, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Trying Heroin Once or Twice (II) among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002

(I) Demographic Characteristic	Try Heroin Once or Twice ¹	(II) Drug Use	Try Heroin Once or Twice ¹	
	Great Risk		Great Risk	Moderate/Slight/No Risk
TOTAL	108	PAST MONTH		
AGE		Any Illicit Drug ²	55	38
12-13	62	Marijuana	48	29
14-15	65	Any Illicit Drug Other Than Marijuana ²	40	30
16-17	55	Cigarettes	59	41
GENDER		Alcohol	66	45
Male	77	Binge Alcohol ³	56	34
Female	74	PAST YEAR		
HISPANIC ORIGIN AND RACE		Any Illicit Drug ²	72	53
Not Hispanic or Latino	98	Marijuana	66	40
White	82	Any Illicit Drug Other Than Marijuana ²	57	45
Black or African American	38	Cigarettes	69	49
American Indian or Alaska Native	7	Alcohol	80	63
Native Hawaiian or Other Pacific Islander	*	LIFETIME		
Asian	27	Any Illicit Drug ²	76	65
Two or More Races	13	Marijuana	71	48
Hispanic or Latino	49	Any Illicit Drug Other Than Marijuana ²	68	56
COUNTY TYPE		Cigarettes	83	59
Large Metro	81	Alcohol	78	67
Small Metro	51			
Nonmetro	44			

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 3.7D Perceived Great Risk of Trying Heroin Once or Twice, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Trying Heroin Once or Twice (II) among Persons Aged 12 to 17: Standard Errors of Percentages, 2002

(I) Demographic Characteristic	Try Heroin Once or Twice ¹	(II) Drug Use	Try Heroin Once or Twice ¹	
	Great Risk		Great Risk	Moderate/Slight/No Risk
TOTAL	0.45	PAST MONTH		
AGE		Any Illicit Drug ²	0.39	0.38
12-13	0.76	Marijuana	0.34	0.29
14-15	0.78	Any Illicit Drug Other Than Marijuana ²	0.28	0.30
16-17	0.69	Cigarettes	0.42	0.40
GENDER		Alcohol	0.46	0.45
Male	0.62	Binge Alcohol ³	0.39	0.33
Female	0.63	PAST YEAR		
HISPANIC ORIGIN AND RACE		Any Illicit Drug ²	0.50	0.52
Not Hispanic or Latino	0.48	Marijuana	0.47	0.40
White	0.54	Any Illicit Drug Other Than Marijuana ²	0.40	0.45
Black or African American	1.09	Cigarettes	0.49	0.49
American Indian or Alaska Native	3.93	Alcohol	0.56	0.62
Native Hawaiian or Other Pacific Islander	*	LIFETIME		
Asian	2.80	Any Illicit Drug ²	0.53	0.65
Two or More Races	3.39	Marijuana	0.50	0.48
Hispanic or Latino	1.27	Any Illicit Drug Other Than Marijuana ²	0.48	0.56
COUNTY TYPE		Cigarettes	0.59	0.59
Large Metro	0.66	Alcohol	0.55	0.67
Small Metro	0.70			
Nonmetro	0.91			

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 3.8C Perceived Great Risk of Using Heroin Once or Twice a Week, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Using Heroin Once or Twice a Week (II) among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002

(I) Demographic Characteristic	Use Heroin Once or Twice a Week ¹	(II) Drug Use	Use Heroin Once or Twice a Week ¹	
	Great Risk		Great Risk	Moderate/Slight/No Risk
TOTAL	82	PAST MONTH		
AGE		Any Illicit Drug ²	62	25
12-13	57	Marijuana	53	18
14-15	49	Any Illicit Drug Other Than Marijuana ²	46	20
16-17	35	Cigarettes	66	24
GENDER		Alcohol	75	26
Male	58	Binge Alcohol ³	62	21
Female	57	PAST YEAR		
HISPANIC ORIGIN AND RACE		Any Illicit Drug ²	83	33
Not Hispanic or Latino	73	Marijuana	74	24
White	54	Any Illicit Drug Other Than Marijuana ²	66	29
Black or African American	33	Cigarettes	78	30
American Indian or Alaska Native	7	Alcohol	91	38
Native Hawaiian or Other Pacific Islander	*	LIFETIME		
Asian	25	Any Illicit Drug ²	92	40
Two or More Races	9	Marijuana	81	29
Hispanic or Latino	39	Any Illicit Drug Other Than Marijuana ²	79	36
COUNTY TYPE		Cigarettes	98	37
Large Metro	66	Alcohol	93	42
Small Metro	40			
Nonmetro	31			

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 3.8D Perceived Great Risk of Using Heroin Once or Twice a Week, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Using Heroin Once or Twice a Week (II) among Persons Aged 12 to 17: Standard Errors of Percentages, 2002

(I) Demographic Characteristic	Use Heroin Once or Twice a Week ¹	(II) Drug Use	Use Heroin Once or Twice a Week ¹	
	Great Risk		Great Risk	Moderate/Slight/No Risk
TOTAL	0.34	PAST MONTH		
AGE		Any Illicit Drug ²	0.31	0.58
12-13	0.70	Marijuana	0.26	0.42
14-15	0.59	Any Illicit Drug Other Than Marijuana ²	0.23	0.47
16-17	0.45	Cigarettes	0.33	0.56
GENDER		Alcohol	0.37	0.61
Male	0.47	Binge Alcohol ³	0.31	0.49
Female	0.48	PAST YEAR		
HISPANIC ORIGIN AND RACE		Any Illicit Drug ²	0.41	0.77
Not Hispanic or Latino	0.36	Marijuana	0.37	0.56
White	0.35	Any Illicit Drug Other Than Marijuana ²	0.33	0.67
Black or African American	0.93	Cigarettes	0.39	0.70
American Indian or Alaska Native	3.93	Alcohol	0.46	0.90
Native Hawaiian or Other Pacific Islander	*	LIFETIME		
Asian	2.58	Any Illicit Drug ²	0.46	0.93
Two or More Races	2.29	Marijuana	0.40	0.69
Hispanic or Latino	1.02	Any Illicit Drug Other Than Marijuana ²	0.40	0.85
COUNTY TYPE		Cigarettes	0.49	0.87
Large Metro	0.54	Alcohol	0.46	0.98
Small Metro	0.55			
Nonmetro	0.64			

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 3.9C Perceived Great Risk of Trying LSD Once or Twice, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Trying LSD Once or Twice (II) among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002

(I) Demographic Characteristic	Try LSD Once or Twice ¹	(II) Drug Use	Try LSD Once or Twice ¹	
	Great Risk		Great Risk	Moderate/Slight/No Risk
TOTAL	109	PAST MONTH		
AGE		Any Illicit Drug ²	44	51
12-13	60	Marijuana	37	42
14-15	67	Any Illicit Drug Other Than Marijuana ²	32	39
16-17	56	Cigarettes	52	50
GENDER		Alcohol	57	57
Male	78	Binge Alcohol ³	46	48
Female	73	PAST YEAR		
HISPANIC ORIGIN AND RACE		Any Illicit Drug ²	62	65
Not Hispanic or Latino	98	Marijuana	56	55
White	79	Any Illicit Drug Other Than Marijuana ²	48	57
Black or African American	39	Cigarettes	62	59
American Indian or Alaska Native	9	Alcohol	70	69
Native Hawaiian or Other Pacific Islander	*	LIFETIME		
Asian	26	Any Illicit Drug ²	69	70
Two or More Races	14	Marijuana	63	60
Hispanic or Latino	48	Any Illicit Drug Other Than Marijuana ²	58	65
COUNTY TYPE		Cigarettes	77	68
Large Metro	85	Alcohol	73	72
Small Metro	51			
Nonmetro	43			

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 3.9D Perceived Great Risk of Trying LSD Once or Twice, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Trying LSD Once or Twice (II) among Persons Aged 12 to 17: Standard Errors of Percentages, 2002

(I) Demographic Characteristic	Try LSD Once or Twice ¹	(II) Drug Use	Try LSD Once or Twice ¹	
	Great Risk		Great Risk	Moderate/Slight/No Risk
TOTAL	0.46	PAST MONTH		
AGE		Any Illicit Drug ²	0.35	0.45
12-13	0.77	Marijuana	0.30	0.37
14-15	0.83	Any Illicit Drug Other Than Marijuana ²	0.26	0.34
16-17	0.71	Cigarettes	0.41	0.44
GENDER		Alcohol	0.45	0.50
Male	0.64	Binge Alcohol ³	0.37	0.42
Female	0.63	PAST YEAR		
HISPANIC ORIGIN AND RACE		Any Illicit Drug ²	0.49	0.58
Not Hispanic or Latino	0.49	Marijuana	0.45	0.49
White	0.53	Any Illicit Drug Other Than Marijuana ²	0.38	0.51
Black or African American	1.13	Cigarettes	0.50	0.52
American Indian or Alaska Native	4.92	Alcohol	0.56	0.61
Native Hawaiian or Other Pacific Islander	*	LIFETIME		
Asian	2.79	Any Illicit Drug ²	0.55	0.62
Two or More Races	3.71	Marijuana	0.50	0.53
Hispanic or Latino	1.26	Any Illicit Drug Other Than Marijuana ²	0.47	0.58
COUNTY TYPE		Cigarettes	0.62	0.60
Large Metro	0.72	Alcohol	0.58	0.64
Small Metro	0.72			
Nonmetro	0.90			

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 3.10C Perceived Great Risk of Using LSD Once or Twice a Week, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Using LSD Once or Twice a Week (II) among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002

(I) Demographic Characteristic	Use LSD Once or Twice a Week ¹	(II) Drug Use	Use LSD Once or Twice a Week ¹	
	Great Risk		Great Risk	Moderate/Slight/No Risk
TOTAL	93	PAST MONTH		
AGE		Any Illicit Drug ²	56	37
12-13	56	Marijuana	47	30
14-15	55	Any Illicit Drug Other Than Marijuana ²	41	29
16-17	47	Cigarettes	61	36
GENDER		Alcohol	69	39
Male	63	Binge Alcohol ³	54	35
Female	66	PAST YEAR		
HISPANIC ORIGIN AND RACE		Any Illicit Drug ²	74	47
Not Hispanic or Latino	79	Marijuana	65	39
White	61	Any Illicit Drug Other Than Marijuana ²	59	41
Black or African American	34	Cigarettes	73	43
American Indian or Alaska Native	9	Alcohol	85	49
Native Hawaiian or Other Pacific Islander	*	LIFETIME		
Asian	25	Any Illicit Drug ²	82	52
Two or More Races	10	Marijuana	72	43
Hispanic or Latino	45	Any Illicit Drug Other Than Marijuana ²	73	47
COUNTY TYPE		Cigarettes	93	48
Large Metro	72	Alcohol	89	51
Small Metro	49			
Nonmetro	35			

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 3.10D Perceived Great Risk of Using LSD Once or Twice a Week, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Using LSD Once or Twice a Week (II) among Persons Aged 12 to 17: Standard Errors of Percentages, 2002

(I) Demographic Characteristic	Use LSD Once or Twice a Week ¹	(II) Drug Use	Use LSD Once or Twice a Week ¹	
	Great Risk		Great Risk	Moderate/Slight/No Risk
TOTAL	0.39	PAST MONTH		
AGE		Any Illicit Drug ²	0.31	0.65
12-13	0.71	Marijuana	0.26	0.53
14-15	0.68	Any Illicit Drug Other Than Marijuana ²	0.22	0.51
16-17	0.60	Cigarettes	0.34	0.64
GENDER		Alcohol	0.38	0.69
Male	0.52	Binge Alcohol ³	0.30	0.62
Female	0.57	PAST YEAR		
HISPANIC ORIGIN AND RACE		Any Illicit Drug ²	0.41	0.83
Not Hispanic or Latino	0.39	Marijuana	0.36	0.70
White	0.41	Any Illicit Drug Other Than Marijuana ²	0.33	0.73
Black or African American	0.98	Cigarettes	0.40	0.76
American Indian or Alaska Native	4.80	Alcohol	0.47	0.87
Native Hawaiian or Other Pacific Islander	*	LIFETIME		
Asian	2.70	Any Illicit Drug ²	0.45	0.92
Two or More Races	2.75	Marijuana	0.40	0.76
Hispanic or Latino	1.19	Any Illicit Drug Other Than Marijuana ²	0.40	0.83
COUNTY TYPE		Cigarettes	0.51	0.85
Large Metro	0.60	Alcohol	0.49	0.90
Small Metro	0.69			
Nonmetro	0.73			

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 3.11C Perceived Great Risk of Having 4 or 5 Drinks of an Alcoholic Beverage Nearly Every Day, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Having 4 or 5 Drinks of an Alcoholic Beverage Nearly Every Day (II) among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002

(I) Demographic Characteristic	Have 4 or 5 Drinks of Alcoholic Beverage Nearly Every Day ¹	(II) Drug Use	Have 4 or 5 Drinks of Alcoholic Beverage Nearly Every Day ¹	
	Great Risk		Great Risk	Moderate/Slight/No Risk
TOTAL	118	PAST MONTH		
AGE		Any Illicit Drug ²	45	47
12-13	60	Marijuana	36	41
14-15	58	Any Illicit Drug Other Than Marijuana ²	34	36
16-17	65	Cigarettes	48	52
GENDER		Alcohol	55	57
Male	81	Binge Alcohol ³	39	50
Female	72	PAST YEAR		
HISPANIC ORIGIN AND RACE		Any Illicit Drug ²	64	59
Not Hispanic or Latino	102	Marijuana	54	53
White	83	Any Illicit Drug Other Than Marijuana ²	53	50
Black or African American	39	Cigarettes	58	60
American Indian or Alaska Native	9	Alcohol	75	63
Native Hawaiian or Other Pacific Islander	*	LIFETIME		
Asian	25	Any Illicit Drug ²	70	65
Two or More Races	11	Marijuana	60	59
Hispanic or Latino	53	Any Illicit Drug Other Than Marijuana ²	61	57
COUNTY TYPE		Cigarettes	74	64
Large Metro	89	Alcohol	80	67
Small Metro	61			
Nonmetro	44			

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 3.11D Perceived Great Risk of Having 4 or 5 Drinks of an Alcoholic Beverage Nearly Every Day, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Having 4 or 5 Drinks of an Alcoholic Beverage Nearly Every Day (II) among Persons Aged 12 to 17: Standard Errors of Percentages, 2002

(I) Demographic Characteristic	Have 4 or 5 Drinks of Alcoholic Beverage Nearly Every Day ¹	(II) Drug Use	Have 4 or 5 Drinks of Alcoholic Beverage Nearly Every Day ¹	
	Great Risk		Great Risk	Moderate/Slight/No Risk
TOTAL	0.48	PAST MONTH		
AGE		Any Illicit Drug ²	0.29	0.51
12-13	0.73	Marijuana	0.24	0.44
14-15	0.68	Any Illicit Drug Other Than Marijuana ²	0.22	0.39
16-17	0.82	Cigarettes	0.31	0.56
GENDER		Alcohol	0.36	0.61
Male	0.65	Binge Alcohol ³	0.26	0.54
Female	0.60	PAST YEAR		
HISPANIC ORIGIN AND RACE		Any Illicit Drug ²	0.42	0.64
Not Hispanic or Latino	0.49	Marijuana	0.35	0.57
White	0.54	Any Illicit Drug Other Than Marijuana ²	0.35	0.54
Black or African American	1.10	Cigarettes	0.38	0.64
American Indian or Alaska Native	4.93	Alcohol	0.49	0.68
Native Hawaiian or Other Pacific Islander	*	LIFETIME		
Asian	2.49	Any Illicit Drug ²	0.46	0.70
Two or More Races	2.88	Marijuana	0.40	0.64
Hispanic or Latino	1.36	Any Illicit Drug Other Than Marijuana ²	0.40	0.62
COUNTY TYPE		Cigarettes	0.49	0.69
Large Metro	0.73	Alcohol	0.52	0.72
Small Metro	0.83			
Nonmetro	0.89			

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 3.12C Perceived Great Risk of Having 5 or More Drinks of an Alcoholic Beverage Once or Twice a Week, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Having 5 or More Drinks of an Alcoholic Beverage Once or Twice a Week (II) among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002

(I) Demographic Characteristic	Have 5 or More Drinks of Alcoholic Beverage Once or Twice a Week ¹	(II) Drug Use	Have 5 or More Drinks of Alcoholic Beverage Once or Twice a Week ¹	
	Great Risk		Great Risk	Moderate/Slight/No Risk
TOTAL	106	PAST MONTH		
AGE		Any Illicit Drug ²	36	58
12-13	62	Marijuana	29	49
14-15	62	Any Illicit Drug Other Than Marijuana ²	25	43
16-17	57	Cigarettes	38	64
GENDER		Alcohol	41	68
Male	71	Binge Alcohol ³	28	60
Female	73	PAST YEAR		
HISPANIC ORIGIN AND RACE		Any Illicit Drug ²	50	77
Not Hispanic or Latino	96	Marijuana	41	67
White	78	Any Illicit Drug Other Than Marijuana ²	37	63
Black or African American	35	Cigarettes	47	73
American Indian or Alaska Native	8	Alcohol	59	82
Native Hawaiian or Other Pacific Islander	*	LIFETIME		
Asian	28	Any Illicit Drug ²	59	84
Two or More Races	13	Marijuana	49	73
Hispanic or Latino	48	Any Illicit Drug Other Than Marijuana ²	50	70
COUNTY TYPE		Cigarettes	62	84
Large Metro	83	Alcohol	62	87
Small Metro	59			
Nonmetro	44			

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 3.12D Perceived Great Risk of Having 5 or More Drinks of an Alcoholic Beverage Once or Twice a Week, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Having 5 or More Drinks of an Alcoholic Beverage Once or Twice a Week (II) among Persons Aged 12 to 17: Standard Errors of Percentages, 2002

(I) Demographic Characteristic	Have 5 or More Drinks of Alcoholic Beverage Once or Twice a Week ¹	(II) Drug Use	Have 5 or More Drinks of Alcoholic Beverage Once or Twice a Week ¹	
	Great Risk		Great Risk	Moderate/Slight/No Risk
TOTAL	0.43	PAST MONTH		
AGE		Any Illicit Drug ²	0.39	0.38
12-13	0.76	Marijuana	0.31	0.32
14-15	0.74	Any Illicit Drug Other Than Marijuana ²	0.26	0.29
16-17	0.72	Cigarettes	0.40	0.42
GENDER		Alcohol	0.44	0.45
Male	0.56	Binge Alcohol ³	0.30	0.39
Female	0.61	PAST YEAR		
HISPANIC ORIGIN AND RACE		Any Illicit Drug ²	0.53	0.51
Not Hispanic or Latino	0.46	Marijuana	0.44	0.44
White	0.51	Any Illicit Drug Other Than Marijuana ²	0.40	0.41
Black or African American	0.98	Cigarettes	0.50	0.48
American Indian or Alaska Native	4.23	Alcohol	0.62	0.54
Native Hawaiian or Other Pacific Islander	*	LIFETIME		
Asian	2.83	Any Illicit Drug ²	0.63	0.55
Two or More Races	3.40	Marijuana	0.52	0.48
Hispanic or Latino	1.24	Any Illicit Drug Other Than Marijuana ²	0.54	0.46
COUNTY TYPE		Cigarettes	0.66	0.55
Large Metro	0.67	Alcohol	0.66	0.57
Small Metro	0.80			
Nonmetro	0.90			

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.